

Genome-wide association identifies *OBFC1* as a locus involved in human leukocyte telomere biology

Daniel Levy^{a,b,c,1}, Susan L. Neuhausen^{d,2}, Steven C. Hunt^{e,2}, Masayuki Kimura^{f,2}, Shih-Jen Hwang^{a,b,c,2}, Wei Chen^{g,2}, Joshua C. Bis^{h,i,2}, Annette L. Fitzpatrick^{j,2}, Erin Smith^k, Andrew D. Johnson^{a,b,c}, Jeffrey P. Gardner^f, Sathanur R. Srinivasan^g, Nicholas Schork^k, Jerome I. Rotter^l, Utz Herbig^f, Bruce M. Psaty^{h,i,m}, Malinee Sastrasinh^f, Sarah S. Murray^k, Ramachandran S. Vasan^{a,b,c}, Michael A. Provinceⁿ, Nicole L. Glazer^{h,i}, Xiaobin Lu^f, Xiaojian Cao^f, Richard Kronmal^o, Massimo Mangino^p, Nicole Soranzo^{p,q}, Tim D. Spector^p, Gerald S. Berenson^g, and Abraham Aviv^{f,1}

^aNational Heart, Lung, and Blood Institute's Framingham Heart Study, Framingham, MA; ^bCenter for Population Studies, National Heart, Lung, and Blood Institute, Bethesda, MD; ^cDivisions of Cardiology and Epidemiology, Boston University School of Medicine, Boston, MA 02118; ^dDepartment Population Sciences, The Beckman Research Institute of the City of Hope, Duarte, CA 91010; ^eCardiovascular Genetics Division, University of Utah School of Medicine, Salt Lake City, UT 84112; ^fCenter of Human Development and Aging, New Jersey Medical School, Newark, NJ 07101; ^gBogalusa Heart Study, Tulane University School of Public Health and Tropical Medicine, New Orleans, LA 70112; ^hCardiovascular Health Research Unit and Department of Medicine, University of Washington, Seattle, WA 98105; ⁱGroup Health Research Institute, Group Health, Seattle, WA 98105; ^jDepartments of Epidemiology and Global Health, University of Washington, Seattle, WA 98105; ^kThe Scripps Translational Science Institute and The Scripps Research Institute, San Diego, CA 92037; ^lMedical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, CA 90048; ^mDepartments of Epidemiology and Health Services, University of Washington, Seattle, WA 98105; ⁿDivision of Statistical Genomics, Washington University School of Medicine, St. Louis, MO 63108; ^oDepartment of Biostatistics, University of Washington, Seattle, WA 98105; ^pDepartment of Twin Research and Genetic Epidemiology, King's College, London SE1 7EH, United Kingdom; and ^qWellcome Trust Sanger Institute, Hinxton CB10 1HH, United Kingdom

Edited by Charles Kooperberg, Fred Hutchinson Cancer Research Center, Seattle, WA, and accepted by the Editorial Board March 31, 2010 (received for review October 6, 2009)

Telomeres are engaged in a host of cellular functions, and their length is regulated by multiple genes. Telomere shortening, in the course of somatic cell replication, ultimately leads to replicative senescence. In humans, rare mutations in genes that regulate telomere length have been identified in monogenic diseases such as dyskeratosis congenita and idiopathic pulmonary fibrosis, which are associated with shortened leukocyte telomere length (LTL) and increased risk for aplastic anemia. Shortened LTL is observed in a host of aging-related complex genetic diseases and is associated with diminished survival in the elderly. We report results of a genome-wide association study of LTL in a consortium of four observational studies ($n = 3,417$ participants with LTL and genome-wide genotyping). SNPs in the regions of the oligonucleotide/oligosaccharide-binding folds containing one gene (*OBFC1*; rs4387287; $P = 3.9 \times 10^{-9}$) and chemokine (C-X-C motif) receptor 4 gene (*CXCR4*; rs4452212; $P = 2.9 \times 10^{-8}$) were associated with LTL at a genome-wide significance level ($P < 5 \times 10^{-8}$). We attempted replication of the top SNPs at these loci through de novo genotyping of 1,893 additional individuals and in silico lookup in another observational study ($n = 2,876$), and we confirmed the association findings for *OBFC1* but not *CXCR4*. In addition, we confirmed the telomerase RNA component (*TERC*) as a gene associated with LTL ($P = 1.1 \times 10^{-5}$). The identification of *OBFC1* through genome-wide association as a locus for interindividual variation in LTL in the general population advances the understanding of telomere biology in humans and may provide insights into aging-related disorders linked to altered LTL dynamics.

genes | aging | *CXCR4* | *TERC*

Leukocyte telomere length (LTL) undergoes progressive attrition with age and is relatively short in aging-related diseases, most notably atherosclerosis (1, 2). LTL displays high interindividual variation at birth (3, 4) and thereafter (5–7), and it is longer in women than in men (8–11) and longer in African Americans than in whites of European descent (10). Although controversy existed regarding the relation of LTL to survival in the elderly, recent studies in same-sex elderly twins have found that the twin with shorter LTL was more likely to die before the twin with longer LTL (12, 13).

LTL is a complex genetic trait; it is heritable (6, 10, 11, 14, 15) and modified by paternal age at conception (5, 16–18). In addition, several environmental factors, including smoking (7, 8), sedentary living (19), obesity (7, 10, 20), low socio-economic status (21), and unhealthy lifestyle (9), are associated with shorter LTL. Shortened LTL has been observed in rare syndromes, including idiopathic aplastic anemia, idiopathic pulmonary fibrosis, and dyskeratosis

congenita (autosomal dominant and X-linked forms) (22–25). These rare syndromes arise from mutations in genes that encode the catalytic (protein) subunit of telomerase reverse transcriptase (*TERT*), the telomerase RNA component (*TERC*), which is the template for the synthesis of telomere repeats, and dyskerin (*DKC1*), a telomerase regulatory protein that binds *TERC*.

Genome-wide screens and genetic mapping have identified more than 150 genes in yeast that are potentially involved in telomere regulation (26, 27). A number of suggestive loci for LTL in humans have been identified through linkage analyses (14, 15, 28) and genome-wide association (GWA) analysis (29). To date, however, these approaches have not identified loci other than *TERC* (30) that are associated with LTL in the general population at a genome-wide level of significance. To identify genetic variants associated with LTL, we established a consortium of four population-based cohorts in which LTL was measured and genome-wide genotyping was performed. We report results of the meta-analysis of GWA of LTL coupled with de novo genotyping and in silico replication of SNPs at the top loci in our GWA study.

Results

Clinical Characteristics. The sample size of individuals with genome-wide genotyping and LTL measurement was 3,417. Table 1 provides the clinical characteristics of participants in the four GWA cohorts: the Framingham Heart Study (31), the Family Heart Study (32), the Cardiovascular Health Study (33, 34), and the Bogalusa Heart Study (35).

GWA Results. Results of the meta-analysis of the GWA of LTL among the four cohorts are presented in Table 2 for SNPs with

Author contributions: D.L., S.C.H., T.D.S., and A.A. designed research; S.L.N., M.K., A.L.F., J.P.G., S.R.S., J.I.R., M.S., S.S.M., X.L., X.C., N. Soranzo, and T.D.S. performed research; R.S.V., M.A.P., G.S.B., and A.A. contributed new reagents/analytic tools; S.C.H., S.-J.H., W.C., J.C.B., E.S., A.D.J., N. Schork, B.M.P., N.L.G., R.K., and M.M. analyzed data; and D.L., S.C.H., N. Schork, U.H., and A.A. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. C.K. is a guest editor invited by the Editorial Board.

Freely available online through the PNAS open access option.

¹To whom correspondence may be addressed. E-mail: avivab@umdnj.edu or Levyd@nih.gov.

²S.L.N., S.C.H., M.K., S.-J.H., W.C., J.C.B., and A.L.F. contributed equally to this work.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.0911494107/-DCSupplemental.

Table 1. Clinical characteristics of the genome-wide association study cohorts

Parameter	Framingham Heart Study	Family Heart Study	Cardiovascular Health Study	Bogalusa Heart Study
Number	1,146	877	1,061	333
Women (%)	51	51	62	42
Age in years (range)	59 (33–86)	62 (30–93)	75 (67–95)	35 (20–48)*
BMI (kg/m ²)	28.0 ± 5.0	29.1 ± 5.6	26.6 ± 4.4	28.0 ± 6.7*

BMI, body mass index ± SD.

*These values include two timepoints for each individual.

$P < 5 \times 10^{-7}$. For the SNPs reported in Table 2, stratified results are provided in Table S1 for the individual cohorts and Table S2 for men and women. In the overall meta-analysis (Table 2), genome-wide significant associations were observed for two regions: the oligonucleotide/oligosaccharide-binding fold containing one gene

(*OBFC1*; lowest $P = 3.9 \times 10^{-9}$ for rs4387287) and the chemokine (C-X-C motif) receptor 4 gene (*CXCR4*; lowest $P = 2.9 \times 10^{-8}$ for rs4452212). Fig. 1, generated with SNAP (36) (<http://www.broadinstitute.org/mpg/snap/>), displays associations in the regions of *OBFC1* and *CXCR4*. Mean LTL values by genotype of the top

Table 2. Meta-analysis of genome-wide association results for all SNPs with $P < 5 \times 10^{-7}$

SNP	Chr	Position	Closest gene	Coded allele	Noncoded allele	Coded allele frequency	β	SE	P
rs4387287	10	105667887	OBFC1	A	C	0.08	0.12	0.02	3.87E-09
rs9420907	10	105666455	OBFC1	A	C	0.91	-0.11	0.02	2.01E-08
rs9419958	10	105665936	OBFC1	T	C	0.08	0.11	0.02	2.08E-08
rs2487999	10	105649816	OBFC1	T	C	0.05	0.13	0.02	2.69E-08
rs4452212	2	136732461	CXCR4	A	G	0.65	-0.08	0.01	2.94E-08
rs4072435	2	136730371	CXCR4	T	C	0.65	-0.08	0.01	2.96E-08
rs13024450	2	136731081	CXCR4	T	C	0.65	-0.08	0.01	2.96E-08
rs13018756	2	136724705	CXCR4	T	C	0.58	-0.08	0.01	2.99E-08
rs13004902	2	136744150	CXCR4	A	G	0.36	0.08	0.01	3.71E-08
rs12691881	2	136746138	CXCR4	A	G	0.35	0.08	0.01	3.76E-08
rs10883943	10	105641406	OBFC1	T	G	0.58	-0.08	0.01	6.09E-08
rs10786775	10	105647306	OBFC1	C	G	0.95	-0.12	0.02	6.13E-08
rs10786774	10	105634313	OBFC1	C	G	0.99	-0.12	0.02	6.29E-08
rs4918068	10	105634726	OBFC1	T	G	0.05	0.12	0.02	6.33E-08
rs11598840	10	105635171	OBFC1	A	G	0.05	0.12	0.02	6.43E-08
rs10883942	10	105641376	OBFC1	T	C	0.42	0.08	0.01	6.49E-08
rs11191849	10	105640870	OBFC1	A	T	0.59	-0.08	0.01	6.50E-08
rs11191848	10	105640827	OBFC1	T	C	0.42	0.08	0.01	6.52E-08
rs10883941	10	105640772	OBFC1	T	C	0.58	-0.08	0.01	6.54E-08
rs2488000	10	105661099	OBFC1	T	G	0.05	0.12	0.02	6.81E-08
rs2488001	10	105661122	OBFC1	A	G	0.05	0.12	0.02	6.96E-08
rs911547	10	105629411	OBFC1	A	G	0.90	-0.11	0.02	6.98E-08
rs1265165	10	105661517	OBFC1	T	C	0.05	0.12	0.02	7.01E-08
rs2756116	10	105661794	OBFC1	T	C	0.05	0.12	0.02	7.05E-08
rs11591710	10	105677622	OBFC1	A	C	0.92	-0.11	0.02	7.87E-08
rs2736428	6	31951903	SLC44A4	T	C	0.29	0.08	0.01	8.59E-08
rs10221893	2	136730076	CXCR4	T	C	0.63	-0.08	0.02	1.12E-07
rs6430612	2	136722668	CXCR4	T	C	0.65	-0.08	0.02	1.15E-07
rs1975174	19	22307091	ZNF676	T	G	0.47	0.07	0.01	1.20E-07
rs9325507	10	105635612	OBFC1	T	C	0.58	-0.07	0.01	1.39E-07
rs7259376	19	22299545	LOC148198	A	G	0.53	-0.07	0.01	1.44E-07
rs11598018	10	105651305	OBFC1	A	C	0.58	-0.07	0.01	1.44E-07
rs12765878	10	105659612	OBFC1	T	C	0.43	0.07	0.01	1.45E-07
rs4954585	2	136714864	CXCR4	T	C	0.75	-0.08	0.02	1.45E-07
rs2273698	10	105639148	OBFC1	A	G	0.58	-0.07	0.01	1.54E-07
rs3814220	10	105637290	OBFC1	A	G	0.43	0.07	0.01	1.55E-07
rs8081000	17	39070759	MEOX1	A	G	0.49	0.08	0.01	1.59E-07
rs351089	4	132636028	PCDH10	T	C	0.99	-0.41	0.08	2.16E-07
rs11668269	19	22305800	LOC148198	A	G	0.47	0.07	0.01	2.33E-07
rs808373	19	22261073	ZNF676	C	G	0.54	-0.07	0.01	2.37E-07
rs8111537	19	22327385	LOC148198	C	G	0.56	-0.07	0.01	3.06E-07
rs1980653	10	105644154	OBFC1	A	G	0.43	0.07	0.01	3.09E-07
rs11191841	10	105629601	OBFC1	T	C	0.42	0.07	0.01	3.11E-07
rs7100920	10	105630968	OBFC1	T	C	0.58	-0.07	0.01	3.25E-07




Fig. 1. Locus-specific association maps for LTL. Locus-specific ($-\log_{10} P$ values) maps in the meta-analysis of LTL for the *OBFC1* (Upper) and *CXCR4* (Lower) loci. The *OBFC1* locus is represented by sentinel SNP rs4387287, and the *CXCR4* locus is represented by rs4452212. For each locus, the sentinel SNP (lowest P value) is depicted by a red diamond, and pairwise linkage disequilibrium in the HapMap CEU with the sentinel SNP is indicated by the color scale. Superimposed on the plot are gene locations (green) and recombination rates (blue).

OBFC1 and *CXCR4* SNPs are provided in Table S3. *OBFC1* was associated with LTL both in men and women (Table S2). In contrast, *CXCR4* was more strongly associated with LTL in women than in men (Wald test for sex difference; $P < 0.0001$). Table 3 displays gene regions associated with LTL in sex-stratified analyses at stratum-specific $P < 5 \times 10^{-7}$ (Wald test for sex differences; $P < 0.0001$ for all SNPs in Table 3). In men, a genome-wide significant association was observed in the region of *DDX18* ($P = 8.2 \times 10^{-9}$ for rs6712766); suggestive association was present for *CCBE1*, *CCDC93*, and *PODXL*. In women, genome-wide significant association was observed in the region of *CNTNAP5* ($P = 3.8 \times 10^{-8}$ for rs10189889); suggestive signals were found for *TNFSF4*, *VPS13C*, and *CXCR4*.

We examined GWA study results for SNPs in the regions of *TERC* and *TERT*, two genes implicated in monogenic diseases associated with shortened LTL. We found evidence of association in the region of *TERC* (rs3772190; $P = 1.13 \times 10^{-5}$) (Table S4); The *TERC* locus was recently reported to be associated with LTL in a meta-analysis that included the TwinsU.K. data (30). Meta-analysis of GWA results for the TwinsU.K. top SNP (rs12696304) compared with those from our four cohorts yielded a P value of 5.5×10^{-8} . We did not detect a signal in the region of *TERT*. We additionally examined associations for SNPs in the region of 18q12.2 and other SNPs recently reported to be associated with LTL (29), but we found no association signal (lowest $P = 0.02$) (Table S5).

Replication Through de Novo Genotyping. Results of the replication genotyping of SNPs in the *OBFC1* and *CXCR4* loci are presented

in Table 4. For the *OBFC1* locus, we genotyped three of the SNPs most strongly associated with LTL (rs4387287, rs9419958, and nonsynonymous rs2487999) in 1,893 additional participants in the Family Heart Study (Table S6). A fourth SNP, rs9420907, was in complete linkage disequilibrium with rs9419958 and therefore, was not genotyped. We found evidence of replication for *OBFC1* in whites ($P = 0.013$ for rs4387287 and $P = 0.027$ for rs9419958) and the combined sample of whites and African Americans ($P = 0.0026$ for rs4387287 and $P = 0.0032$ for rs9419958). For the *CXCR4* locus, we genotyped rs4452212 and rs4954585 in additional samples from the Family Heart Study but were unable to replicate the GWA findings (Table 4).

Replication Through in Silico Lookup. Replication of SNPs at our top loci also was undertaken through lookup of results within the LTL GWA of women from the TwinsU.K. Study (30), which had genome-wide genotyping and LTL data for 2,876 women. Six SNPs from the combined sex analysis of Table 2 (rs351089, rs4452212, rs2736428, rs1975174, rs8081000, and rs4387287) were looked up in the TwinsU.K. Study and meta-analyzed in conjunction with the results from the four GWA cohorts; the total sample size was 6,293. The *OBFC1* locus replicated (rs4387287, $P = 8.9 \times 10^{-4}$ in TwinsU.K. Study; $P = 2.3 \times 10^{-11}$ in the meta-analysis of our GWA plus TwinsU.K. Study). Table 5 summarizes these results.

Discussion

Our GWA meta-analysis identified genome-wide significant associations of LTL with regions containing *OBFC1* and *CXCR4*.

Table 3. Meta-analysis of genome-wide association results for men and women for SNPs with $P < 5 \times 10^{-7}$ in either sex

SNP	Chr	Position	Closest gene	Coded allele	Coded allele frequency	Meta-GWA study results for women			Meta-GWA study results for men			Wald test*	
						β	SE	P	β	SE	P	T value	P
rs6712766	2	118328259	DDX18	A	0.07	-0.04	0.04	3.66E-01	-0.21	0.04	8.22E-09	34.12	<0.0001
rs1791285	18	55517163	CCBE1	T	0.80	0.03	0.03	3.70E-01	-0.16	0.03	1.25E-07	28.78	<0.0001
rs13432259	2	118361546	CCDC93	A	0.04	-0.06	0.04	1.27E-01	-0.20	0.04	2.04E-07	29.27	<0.0001
rs12534413	7	131104504	PODXL	T	0.17	0.02	0.03	5.96E-01	-0.15	0.03	2.21E-07	27.17	<0.0001
rs10189889	2	125161934	CNTNAP5	A	0.03	0.34	0.06	3.77E-08	-0.06	0.06	3.47E-01	31.18	<0.0001
rs11583523	1	171429823	TNFSF4	A	0.99	-0.98	0.19	1.61E-07	-0.18	0.16	2.57E-01	28.73	<0.0001
rs9920256	15	59986872	VPS13C	A	0.03	0.37	0.07	2.66E-07	0.04	0.07	6.01E-01	26.79	<0.0001
rs4452212	2	136732461	CXCR4	A	0.65	-0.10	0.02	4.27E-07	-0.05	0.02	9.84E-03	32.15	<0.0001

*Men vs. women.

Our association results for *OBFC1* were replicated through de novo genotyping of 1,893 additional individuals and through silico analysis of 2,876 individuals with existing LTL GWA data; however, we were unable to replicate our findings for *CXCR4*. We also were able to confirm a recently reported association for the *TERC* locus (30). The potential implications of these findings are discussed based on the postulate that LTL dynamics (birth LTL and its age-dependent shortening afterward) mirror telomere dynamics in hematopoietic stem cells (HSCs) from which leukocytes are derived (37).

Theoretical considerations suggest that genes associated with LTL belong to two general categories (i.e., genes directly engaged in telomere maintenance and genes that impact the turnover rate of HSCs). Based on the biology of the genes implicated in our study, we contend that *OBFC1* and *TERC* fall into the former category and *CXCR4* falls into the latter category.

OBFC1 is the human homolog of yeast Stn1 (38, 39), a protein specifically involved in the replication and capping of telomeres (40). Yeast Stn1 regulates synthesis of the telomeric G-rich strand by both protecting terminal telomeric DNA and negatively regulating telomerase action on telomeres (41, 42). In addition, Stn1 coordinates DNA replication of the opposing telomeric C strand. Importantly, it has been recently shown that *OBFC1* interacts and colocalizes with telomeric proteins in human cells, suggesting that *OBFC1* regulates telomere length and/or function in humans (38, 39). Supporting this is the observation that overexpression of an *OBFC1* truncation mutation causes telomere elongation in cancer cells (38).

Interrogation of SNP databases revealed that two of the top *OBFC1* SNPs in our LTL GWA results are nonsynonymous and in tight-linkage disequilibrium (rs2487999, $P = 2.7 \times 10^{-8}$; rs10786775, $P = 6.1 \times 10^{-8}$; pairwise $r^2 = 1.0$). They are predicted by PolyPhen (<http://genetics.bwh.harvard.edu/pph/>) to be benign individually, but the effects of two shared nonsynonymous amino acid substitutions in the coded protein are not known.

Because *OBFC1* is a major gene that is directly engaged in telomere biology, we genotyped the top GWA results for *OBFC1* in participants from the Family Heart Study and conducted in silico analysis with GWA results from the TwinsU.K. Study in an

effort to replicate our findings. We found evidence of replication for *OBFC1* in the genotyped individuals from the Family Heart Study (Table 4) and in the in silico analysis of women from the TwinsU.K. Study (Table 5). In our genotyped replication samples, on average, the difference in LTL across *OBFC1* genotypes of rs4387287 and rs9419958 amounted to 230 bp and 290 bp in whites, respectively (Table 4). In the meta-analysis, the mean LTL difference across genotypes was 400 bp in men and 180 bp in women for rs437287 (Table S3). These differences in telomere length by genotype are substantial, given that the rate of LTL shortening in the general population is about 20–30 bp/year (43).

Of note, the Framingham Heart Study previously reported an association of the *OBFC1* locus (rs3814219; $P = 9.5 \times 10^{-7}$) with brachial-artery basal blood flow, which is a functional index of the vascular endothelium (44). However, the top SNP in the GWA of vascular function was weakly correlated with the top *OBFC1* SNPs in this investigation (rs3814219 and rs4387287, $r^2 = 0.02$; other pairwise r^2 values did not exceed 0.34).

Although we were not able to replicate our *CXCR4* results, several lines of evidence point to a connection between *CXCR4* and LTL-related biology. First, *CXCR4* is a key regulator of the trafficking of neutrophils between the bone marrow and blood (45, 46). Thus, *CXCR4* variants might impact HSC turnover and consequently, LTL. Second, shortened LTL is associated with atherosclerosis (1, 2), an inflammatory disease of the vascular endothelium, marked by a damage-repair feedback loop between the bone marrow (i.e., HSCs) and vascular endothelium. This feedback is mediated by chemotactic signals and endothelial progenitor cells. At its center are chemokines, specifically, *CXCR4* and its cognate ligand *CXCL12* (47, 48). Moreover, the *CXCL12* locus has been found to be associated with myocardial infarction at a genome-wide level of significance (49). Despite these enticing physiological observations, our inability to replicate the association of *CXCR4* with LTL deserves further investigation to determine if the LTL association with *CXCR4* in our meta-analysis is a false-positive result.

Finally, mutations in *TERT*, *TERC*, and *DKC1* have been identified in relatively rare monogenic diseases that are associated with shortened telomeres. A lookup of our association results identified an association signal in the *TERC* locus (rs3772190; $P = 1.1 \times 10^{-5}$); this SNP is in strong linkage disequilibrium with a newly reported GWA finding at this locus (rs12696304, P value in our meta-analysis = 1.6×10^{-4} ; pairwise $r^2 = 0.91$) (Table S4) (30). Combining the results from the TwinsU.K. Study and our study produced a P value of 5.5×10^{-8} , confirming that this locus is associated with LTL.

Given that shortened LTL denotes susceptibility to aging-related diseases, principally atherosclerosis, in the general population, the associations of LTL with *OBFC1* and *TERC* (and potentially, *CXCR4* if replicated by other studies) bolster the tenet that telomere biology is an important pathway in human aging. Although many factors are at work in fashioning human LTL, our findings are arguably a major line of evidence that genes engaged in telomere maintenance (and perhaps, HSC turnover) are directly involved in this process and explain, in part, interindividual variation in LTL in the general population.

Materials and Methods

The LTL Consortium includes cohorts from four studies with genome-wide genotyping and LTL measurements: the Framingham Heart Study, the Family Heart Study, the Cardiovascular Health Study, and the Bogalusa Heart Study (SI Text). The consortium established a consensus on covariate selection a priori and an analytical plan for within-study GWA and meta-analysis of results across the studies. Each study received institutional review-board approval of its research procedures, procedures for DNA collection, and use for genetic research. All participants in each cohort study gave written informed consent for participation in the parent study and for the conduct of genetic research using DNA.

Table 4. Replication results for *OBFC1* and *CXCR4*

<i>OBFC1</i> genotype*	1	2	3	<i>P</i>
Whites of European descent ($n = 1,319$)				
rs4387287 (C/A)	6.80 ± 0.024	6.88 ± 0.038	7.03 ± 0.083	0.013
rs9419958 (C/T)	6.80 ± 0.023	6.90 ± 0.040	7.09 ± 0.112	0.027
rs2487999 (C/T)	6.81 ± 0.023	6.92 ± 0.044	7.08 ± 0.20	0.21
African Americans ($n = 574$)				
rs4387287 (A/C)	7.08 ± 0.042	7.03 ± 0.042	7.01 ± 0.071	0.34
rs9419958 (T/C)	7.10 ± 0.055	7.06 ± 0.042	6.96 ± 0.053	0.062
rs2487999 (C/T)	6.99 ± 0.042	7.11 ± 0.041	7.07 ± 0.073	0.33
Both races ($n = 1,893$)				
rs4387287 (C/A)	6.89 ± 0.028	6.96 ± 0.029	7.04 ± 0.038	0.0026
rs9419958 (C/T)	6.89 ± 0.025	6.99 ± 0.029	7.06 ± 0.049	0.0032
rs2487999 (C/T)	6.91 ± 0.023	7.03 ± 0.030	7.02 ± 0.067	0.12
<i>CXCR4</i> genotype*				
Whites of European descent ($n = 1,593$)				
rs4452212 (A/G)	6.85 ± 0.031	6.83 ± 0.026	6.83 ± 0.039	0.77
rs4954585 (T/C)	6.83 ± 0.027	6.84 ± 0.025	6.82 ± 0.046	0.86
African Americans ($n = 597$)				
rs4452212 (G/A)	7.06 ± 0.036	7.02 ± 0.049	6.84 ± 0.115	0.098
rs4954585 (C/T)	7.04 ± 0.046	7.09 ± 0.039	6.95 ± 0.065	0.26

Mean LTL (kb) according to genotype.

*Genotype 1 indicates zero copies, genotype 2 indicates one copy, and genotype 3 indicates two copies of the minor allele. The minor allele is listed second next to the rs number for each SNP. Note that the minor alleles may differ by race. P values reflect an additive genetic model.

Table 5. Replication through in silico lookup and meta-analysis with TwinsU.K. Study

SNP	Chr	Position	Nearest gene	Coded allele	Noncoded allele	TwinsU.K. women only (n = 2,876)			Meta-analysis of all studies (n = 6,293)		
						β	SE	P	β	SE	P
rs1975174	19	22307091	ZNF676	T	G	0.02	0.02	2.40E-01	0.05	0.01	2.05E-06
rs2736428	6	31951903	SLC44A4	T	C	0.02	0.02	3.64E-01	0.05	0.01	3.24E-06
rs351089	4	132636028	PCDH10	T	C	-0.06	0.08	4.25E-01	-0.23	0.06	2.86E-05
rs4387287	10	105667887	OBFC1	A	C	0.08	0.03	8.92E-04	0.10	0.02	2.33E-11
rs4452212	2	136732461	CXCR4	A	G	-0.02	0.02	3.31E-01	-0.05	0.01	1.68E-06
rs8081000	17	39070759	MEOX1	A	G	0.00	0.02	9.26E-01	0.04	0.01	1.43E-04

LTL Measurement. LTL was measured by Southern blot analysis of the mean length of terminal restriction fragments as described previously (50, 51). The overlay method was used for the Framingham Heart Study and the Family Heart Study (50), whereas the standard method was used for the Cardiovascular Health Study and the Bogalusa Heart Study (51). These two approaches are highly correlated (50). The laboratory measuring LTL was blind as to the characteristics of DNA donors in all four studies.

Genome-Wide Genotyping. The platforms used for genome-wide genotyping are provided in Table S7. All physical positions use National Center for Biotechnology Information (NCBI) build 36.3 as the reference.

Framingham Heart Study. Genotyping was conducted on 9,274 Framingham Heart Study participants using the Affymetrix 500K mapping array and the Affymetrix 50K gene-focused molecularly imprinted polymer array (SNP Health Association Resource at http://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000007.v2.p1). Individuals were excluded from association analyses when the call rate across all genotyped SNPs was less than 97%. There were 503,551 SNPs available for analysis after exclusions. The genotyped sample fulfilling eligibility criteria with LTL measurements and full covariate information was 1,146 individuals. Researchers imputed the genotype data for 2.5 million SNPs using MACH (version 1.0.15) software for autosomal SNPs. The reference panel for the imputation was the publicly available phased haplotypes from HapMap (release 22, build 26, CEU population). The final genotyping data for analysis are composed of 2,543,887 SNPs in HapMap using allele dosage (0–2) values. Because family members were included within the offspring cohort, significance of the association between SNPs and LTL was tested using the linear mixed-effects regression (LME) model, which takes into consideration random individual effects correlated within pedigree according to kinship relationships.

Family Heart Study. All subjects from the Family Heart Study were genotyped on the Illumina HumMap 550 chip. Of these, 33 (3.3%) were excluded because of technical errors, call rates below 98%, or discrepancies between reported sex and sex-diagnostic markers. There was no significant plate-to-plate variation in allele frequencies. Because the Framingham site of the Family Heart Study was involved in both the Framingham Heart Study and the Family Heart Study, subjects who were in both studies were removed from the Family Heart Study dataset before analysis. For the replication studies, subjects were genotyped after excluding overlap with Framingham subjects and those without LTL measurements or other analysis variables. There were 547,353 SNP markers available for analysis in the Family Heart Study. After exclusions for deviations from Hardy–Weinberg equilibrium ($P < 10^{-6}$), minor allele frequency $< 1\%$, and markers not in the HapMap, 456,293 markers were used as a framework map for imputation. MACH (version 1.0.15) was used to impute up to 2,543,887 SNPs using the publicly available phased haplotypes from HapMap (release 22, build 26, CEU population) as a reference population. A random sample of 200 (100 males and 100 females) unrelated individuals, excluding those with the highest rates of missing genotypes, was used to estimate parameters that were then applied to the remaining subjects. Genotype dosage was output and used in the analysis. An additive linear-regression model using PLINK (<http://pngu.mgh.harvard.edu/~purcell/plink/>) was used for the association analysis of each SNP with LTL. Although some of the 877 subjects were related, the correlation of P values for two chromosomes with and without adjustment for relatedness was over 0.95 with no difference in mean P value. Therefore, no adjustment for relatedness was performed for the analysis of the 877 subjects. For the replication samples, which were family-based, a mixed model was used to account for relatedness separately for whites and African Americans (52).

Cardiovascular Health Study. Genotyping was performed using the Illumina 370CNV BeadChip system on 3,980 Cardiovascular Health Study (CHS) participants who were free of cardiovascular disease at baseline. Participants were excluded if they had a call rate $\leq 95\%$. Because the other cohorts were predominantly white, the African-American participants were excluded from this analysis. To date, genotyping has been successful in 3,291 of the attempted 3,397 white participants. A total of 1,080 samples from individuals of European ancestry were available with both LTL measurements and GWA analysis. The following exclusions were applied to identify a final set of 306,655 autosomal SNPs: call rate $< 97\%$, Hardy–Weinberg Equilibrium $P < 10^{-5}$, more than two duplicate errors or Mendelian inconsistencies (for reference Centre d'Etude du Polymorphisme Humain trios), heterozygote frequency = 0, or SNP not found in HapMap. Imputation was then performed using BAMBAM v0.99 with reference to HapMap CEU using release 22, build 36 with one round of imputations and the default expectation-maximization warm-ups and runs. During the analysis, SNPs were excluded for variance on the allele dosage ≤ 0.01 .

Bogalusa Heart Study. Individuals were genotyped on the Illumina Human610 Genotyping Beadchip (53, 54). Genotypes were called using the BeadStudio clustering algorithm. Samples were filtered based on the 10th and 50th percentile GenCall score and overall call rates ($< 99\%$). Additionally, some related individuals were present in the sample and were identified using Identity-by-Descent measures in PLINK (55). Individuals were filtered such that there were no pairs with π -hat > 0.10 . This resulted in a total of 333 individuals. SNPs were filtered based on call rates ($< 90\%$). Cluster plots for SNPs with call rates between 90% and 95% or with cluster separation scores < 0.30 were manually inspected. By genotyping 29 samples in duplicate (18 known replicates and 11 blind replicates), we observed reproducibility $> 99.99\%$. There were 626,145 genotyped SNPs used in the reproducibility analysis. Imputation was performed with MACH (v. 1.0.16) using phased CEU haplotypes from HapMap Phase 2 (release 22) and 550,798 genotyped SNPs. A random sample of 200 individuals was used to estimate recombination and error rates before imputing the entire sample. This resulted in imputing 2,543,887 SNPs. By masking 0.1% of the genotypes before imputation, we observed allelic error rates of 1.6% and genotypic error rates of 3.1%.

Telomere length was measured at two timepoints for each individual in the Bogalusa Heart Study (BHS). We modeled and tested associations between SNPs and the two LTL measures using a linear mixed model as implemented in the nlme R package (56). A covariance structure for the two LTL measurements was chosen by testing all spatial structures with the model, and the structure that resulted in the lowest Akaike Information Criterion score was used (in this study, the best covariance was exponential). If an SNP had been genotyped, the genotype value (0, 1, or 2) was used as a predictor, whereas if the SNP was imputed, the estimated dosage was used.

GWA Analysis. In each cohort, GWA was conducted for the continuous variable LTL (in kb pairs) using an additive genetic model with adjustment for age, age², sex, cigarette smoking, and body mass index (BMI). GWA was conducted with 2.5 million HapMap SNPs in conjunction with measured genotypes or allele dose when genotypes were imputed. In the Framingham Heart Study where family members were included within the offspring cohort, significance of the association between SNPs and LTL was tested using the linear mixed-effects regression (LME) model, which takes into consideration random individual effects correlated within pedigree according to kinship relationships. Meta-analysis was performed using inverse variance weighting, and a level of statistical significance was established at 5×10^{-8} . A P value between 5×10^{-7} and 5×10^{-8} was considered strongly suggestive of association.

Replication Genotyping. Genotyping was performed using MGB Taqman probe assays from Applied Biosystems. Primer and probe sequences are available on request. Specifically, the reaction mix in a final volume of 5 μ l included 10 ng genomic DNA, 4.5 pmol of each primer, 1.25 pmol of each probe, 1 \times PCR buffer from Qiagen, 2 \times Q solution from Qiagen, 500 pmol dNTP, and 0.15 units of Qiagen DNA polymerase. PCR cycling included 55 cycles of a two-step PCR (95 $^{\circ}$ C for 15 seconds and 60 $^{\circ}$ C for 1 min) after an initial 2 min at 95 $^{\circ}$ C. PCR plates were read on an ABI PRISM 7900 HT instrument for genotype assignment. Duplicates of 22 DNA samples and water controls were genotyped for quality control. The laboratory technician was blinded as to which samples were duplicates or water controls. The order of the DNA samples in the 384-well plates was randomized to ensure balance in study conditions across covariates. Genotyping call rates ranged from 95%

to 99%, and duplicate concordance rates were higher than 99%. For replication analyses, a *P* value of 0.05 divided by the number of SNPs tested for replication was considered statistically significant.

In Silico Replication. The TwinsU.K. Study (29, 30) conducted GWA of LTL measured by the Southern blot method (51). Methods of genome-wide genotyping and GWA have been summarized previously (29). For replication purposes, in silico lookup was carried out in TwinsU.K. women for the top SNPs from the GWA meta-analysis of the four primary cohorts of this investigation. The sample size of TwinsU.K. women for this analysis was 2,876. These results were meta-analyzed with the primary GWA results for these SNPs in our GWA. The total sample size for this combined analysis was 6,293.

- Aviv A (2006) Telomeres and human somatic fitness. *J Gerontol A Biol Sci Med Sci* 61: 871–873.
- Samani NJ, van der Harst P (2008) Biological ageing and cardiovascular disease. *Heart* 94:537–539.
- Okuda K, et al. (2002) Telomere length in the newborn. *Pediatr Res* 52:377–381.
- Akkad A, et al. (2006) Telomere length in small-for-gestational-age babies. *BJOG* 113: 318–323.
- Njajou OT, et al. (2007) Telomere length is paternally inherited and is associated with parental lifespan. *Proc Natl Acad Sci USA* 104:12135–12139.
- Slagboom PE, Droog S, Boomsma DI (1994) Genetic determination of telomere size in humans: A twin study of three age groups. *Am J Hum Genet* 55:876–882.
- Valdes AM, et al. (2005) Obesity, cigarette smoking, and telomere length in women. *Lancet* 366:662–664.
- Nawrot TS, Staessen JA, Gardner JP, Aviv A (2004) Telomere length and possible link to X chromosome. *Lancet* 363:507–510.
- Bekaert S, et al. (2007) Telomere length and cardiovascular risk factors in a middle-aged population free of overt cardiovascular disease. *Aging Cell* 6:639–647.
- Hunt SC, et al. (2008) Leukocyte telomeres are longer in African Americans than in whites: The National Heart, Lung, and Blood Institute Family Heart Study and the Bogalusa Heart Study. *Aging Cell* 7:451–458.
- Jeandros E, et al. (2000) Telomere length inversely correlates with pulse pressure and is highly familial. *Hypertension* 36:195–200.
- Kimura M, et al. (2008) Telomere length and mortality: A study of leukocytes in elderly Danish twins. *Am J Epidemiol* 167:799–806.
- Bakaysa SL, et al. (2007) Telomere length predicts survival independent of genetic influences. *Aging Cell* 6:769–774.
- Andrew T, et al. (2006) Mapping genetic loci that determine leukocyte telomere length in a large sample of unselected female sibling pairs. *Am J Hum Genet* 78:480–486.
- Vasa-Nicotera M, et al. (2005) Mapping of a major locus that determines telomere length in humans. *Am J Hum Genet* 76:147–151.
- Unryn BM, Cook LS, Riabowol KT (2005) Paternal age is positively linked to telomere length of children. *Aging Cell* 4:97–101.
- De Meyer T, et al. (2007) Paternal age at birth is an important determinant of offspring telomere length. *Hum Mol Genet* 16:3097–3102.
- Kimura M, et al. (2008) Offspring's leukocyte telomere length, paternal age, and telomere elongation in sperm. *PLoS Genet* 4:e37.
- Cherkas LF, et al. (2008) The association between physical activity in leisure time and leukocyte telomere length. *Arch Intern Med* 168:154–158.
- Gardner JP, et al. (2005) Rise in insulin resistance is associated with escalated telomere attrition. *Circulation* 111:2171–2177.
- Cherkas LF, et al. (2006) The effects of social status on biological aging as measured by white-blood-cell telomere length. *Aging Cell* 5:361–365.
- Mason PJ, Wilson DB, Bessler M (2005) Dyskeratosis congenita—a disease of dysfunctional telomere maintenance. *Curr Mol Med* 5:159–170.
- Yamaguchi H, et al. (2005) Mutations in TERT, the gene for telomerase reverse transcriptase, in aplastic anemia. *N Engl J Med* 352:1413–1424.
- Tsakiri KD, et al. (2007) Adult-onset pulmonary fibrosis caused by mutations in telomerase. *Proc Natl Acad Sci USA* 104:7552–7557.
- Armanios MY, et al. (2007) Telomerase mutations in families with idiopathic pulmonary fibrosis. *N Engl J Med* 356:1317–1326.
- Askree SH, et al. (2004) A genome-wide screen for *Saccharomyces cerevisiae* deletion mutants that affect telomere length. *Proc Natl Acad Sci USA* 101:8658–8663.
- Gatbonton T, et al. (2006) Telomere length as a quantitative trait: Genome-wide survey and genetic mapping of telomere length-control genes in yeast. *PLoS Genet* 2: e35.
- Mangino M, et al. (2008) A regulatory SNP of the BICD1 gene contributes to telomere length variation in humans. *Hum Mol Genet* 17:2518–2523.
- Mangino M, et al. (2009) A genome-wide association study identifies a novel locus on chromosome 18q12.2 influencing white cell telomere length. *J Med Genet* 46: 451–454.
- Codd V, et al. (2010) Common variants near TERC are associated with mean telomere length. *Nat Genet* 42:197–199.
- Feinleib M, Kannel WB, Garrison RJ, McNamara PM, Castelli WP (1975) The Framingham Offspring Study. Design and preliminary data. *Prev Med* 4:518–525.
- Higgins M, et al. (1996) NHLBI Family Heart Study: Objectives and design. *Am J Epidemiol* 143:1219–1228.
- Fried LP, et al. (1991) The Cardiovascular Health Study: Design and rationale. *Ann Epidemiol* 1:263–276.
- Tell GS, et al. (1993) Recruitment of adults 65 years and older as participants in the Cardiovascular Health Study. *Ann Epidemiol* 3:358–366.
- Berenson GS, et al. (1980) *Cardiovascular Risk Factors in Children—The Early Natural History of Atherosclerosis and Essential Hypertension* (Oxford University Press, New York), pp 47–123.
- Johnson AD, et al. (2008) SNAP: A web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* 24:2938–2939.
- Sidorov I, Kimura M, Yashin A, Aviv A (2009) Leukocyte telomere dynamics and human hematopoietic stem cell kinetics during somatic growth. *Exp Hematol* 37: 514–524.
- Wan M, Qin J, Songyang Z, Liu D (2009) OB fold-containing protein 1 (OBFC1), a human homolog of yeast Stn1, associates with TPP1 and is implicated in telomere length regulation. *J Biol Chem* 284:26725–26731.
- Miyake Y, et al. (2009) RPA-like mammalian Ctc1-Stn1-Ten1 complex binds to single-stranded DNA and protects telomeres independently of the Pot1 pathway. *Mol Cell* 36:193–206.
- Palm W, de Lange T (2008) How shelterin protects mammalian telomeres. *Annu Rev Genet* 42:301–334.
- Puglisi A, Bianchi A, Lemmens L, Damay P, Shore D (2008) Distinct roles for yeast Stn1 in telomere capping and telomerase inhibition. *EMBO J* 27:2328–2339.
- Li S, et al. (2009) Cdk1-dependent phosphorylation of Cdc13 coordinates telomere elongation during cell-cycle progression. *Cell* 136:50–61.
- Aviv A (2008) The epidemiology of human telomeres: Faults and promises. *J Gerontol A Biol Sci Med Sci* 63:979–983.
- Vasan RS, et al. (2007) Genome-wide association of echocardiographic dimensions, brachial artery endothelial function and treadmill exercise responses in the Framingham Heart Study. *BMC Med Genet* 8 (Suppl 1):S2.
- Eash KJ, Means JM, White DW, Link DC (2009) CXCR4 is a key regulator of neutrophil release from the bone marrow under basal and stress granulopoiesis conditions. *Blood* 113:4711–4719.
- Furze RC, Rankin SM (2008) Neutrophil mobilization and clearance in the bone marrow. *Immunology* 125:281–288.
- Zernecke A, Shagdarsuren E, Weber C (2008) Chemokines in atherosclerosis: An update. *Arterioscler Thromb Vasc Biol* 28:1897–1908.
- Bernhagen J, et al. (2007) MIF is a noncognate ligand of CXC chemokine receptors in inflammatory and atherogenic cell recruitment. *Nat Med* 13:587–596.
- Kathiresan S, et al. (2009) Genome-wide association of early-onset myocardial infarction with single nucleotide polymorphisms and copy number variants. *Nat Genet* 41:334–341.
- Vasan RS, et al. (2008) Association of leukocyte telomere length with circulating biomarkers of the renin-angiotensin-aldosterone system: The Framingham Heart Study. *Circulation* 117:1138–1144.
- Benetos A, et al. (2001) Telomere length as an indicator of biological aging: The gender effect and relation with pulse pressure and pulse wave velocity. *Hypertension* 37:381–385.
- Price AL, et al. (2006) Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet* 38:904–909.
- Eberle MA, et al. (2007) Power to detect risk alleles using genome-wide tag SNP panels. *PLoS Genet* 3:1827–1837.
- Keating BJ, et al. (2008) Concept, design and implementation of a cardiovascular gene-centric 50 k SNP array for large-scale genomic association studies. *PLoS One* 3: e3583.
- Purcell S, et al. (2007) PLINK: A tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 81:559–575.
- Pinheiro J, et al. (2009) nlme: Linear and Nonlinear Mixed Effects Models. R package Version 3. Available at <http://cran.r-project.org/web/packages/nlme/citation.html>.